


Claims

1.



in which

R^1 means a monocyclic or bicyclic C_{6-12} aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-4 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br, I,

$C(NH)NH_2$, $C(NH)NHR^4$, $C(NH)NR^4R^{4'}$, $C(NR^4)NH_2$, $C(NR^4)NHR^{4'}$,
 $C(NR^4)NR^4R^{4'}$,

XOH , XOR^4 , $XOCOR^4$, $XOCONHR^4$, $XOCOOR^4$,

$XCOR^4$, $XC(NO)R^4$, $XC(NOR^4)R^{4'}$, $XC(NO(COR^4))R^{4'}$

XCN , $XCOOH$, $XCOOR^4$, $XCONH_2$, $XCONR^4R^{4'}$, $XCONHR^4$, $XCONHOH$,
 $XCONHOR^4$, $XCOSR^4$

XSR^4 , $XSOR^4$, XSO_2R^4 ,

SO_2NH_2 , SO_2NHR^4 , $SO_2NR^4R^{4'}$,

09759360-011604

*Sub
B
cont*

NO_2 , XNH_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$, XNHSO_2R^4 , $\text{XN}(\text{SO}_2\text{R}^4)\text{SO}_2\text{R}^{4'}$,
 $\text{XNR}^4\text{SO}_2\text{R}^{4'}$,

XNHCO^4 , XNHCOOR^4 , XNHCONHR^4 , tetrahydro-2,5-dioxopyrrol-1-yl, 2,5-dihydro-2,5-dioxopyrrol-1-yl, 2,7-dihydro-2,7-dioxoisindol-1-yl, R^4 ,

whereby two substituents at R^1 , if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediylbisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl,

R^2 means a monocyclic or bicyclic C_{6-10} aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-4 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br, I,

XOH , XOR^4 , XOCOR^4 , XOCONHR^4 , XOCOOR^4 ,

XCOR^4 , $\text{XC}(\text{NOH})\text{R}^4$, $\text{XC}(\text{NOR}^4)\text{R}^{4'}$, $\text{XC}(\text{NO}(\text{COR}^4))\text{R}^{4'}$,

XCOOH , XCOOR^4 , XCONH_2 , XCONHR^4 , $\text{XCONR}^4\text{R}^{4'}$, XCONHOH ,

XCONHOR^4 , XCOSR^4 ,

XSR^4 , XSOR^4 , XSO_2R^4 , SO_2NH_2 , SO_2NHR^4 , $\text{SO}_2\text{NR}^4\text{R}^{4'}$,

NO_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$, XNHSO_2R^4 , $\text{XN}(\text{SO}_2\text{R}^4)\text{SO}_2\text{R}^{4'}$,

$\text{XNR}^4\text{SO}_2\text{R}^{4'}$, tetrahydro-2,5-dioxopyrrol-1-yl, 2,5-dihydro-2,5-dioxopyrrol-1-yl, 2,7-dihydro-2,7-dioxoisindol-1-yl, R^4 ,

FOI 0970 09665250

Sub
B
cont

whereby two substituents at R^2 , if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediyl-bisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl,

R^3 means one or two substituents, which form, independently of one another:

hydrogen,

F, Cl, Br, I,

XOH, XOR^4 , $XOCOR^4$, $XOCONHR^4$, $XOCOOR^4$,

$XCOR^4$, $XC(NOHR^4)$, $XC(NOR^4)R^{4'}$, $XC(NO(COR^4))R^{4'}$,

XCN, XCOOH, $XCOOR^4$, $XCONH_2$, $XCONHR^4$, $XCONR^4R^{4'}$, $XCONHOH$,

$XCONHOR^4$, $XCOSR^4$, XSR^4 , $XSOR^4$, XSO_2R^4 , SO_2NH_2 , SO_2NHR^4 ,

$SO_2NR^4R^{4'}$,

NO_2 , XNH_2 , $XNHR^4$, $XNR^4R^{4'}$,

$XNH SO_2R^4$, $XNR^4SO_2R^{4'}$, $XN(SO_2R^4)(SO_2R^{4'})$,

$XNH COR^4$, $XNH COOR^4$, $XNH CONHR^4$, tetrahydro-2,5-

dioxopyrrol-1-yl, 2,5-dihydro-2,5-dioxopyrrol-1-yl,

2,7-dihydro-2,7-dioxoisindol-1-yl, or R^3 can be R^4 ,

whereby two substituents at R^3 , if they are in ortho-position to one another, can be linked to one another

in such a way that they jointly form methanediylbisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl,

R^4 and $R^{4'}$, independently of one another, mean C_{1-4}

perfluoroalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_3 -cycloalkyl, (C_{1-3} alkyl- C_{3-7} cycloalkyl), C_{1-3} alkyl- C_{6-10}

09759360.01601

*Sub
B
cont*

aryl, C_{1-3} alkyl-5 to 10-membered heteroaryl, with 1-4 N, S or O atoms, C_{6-10} aryl or 5- to 10-membered heteroaryl with 1-4 N, S or O atoms, whereby the aryl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F, Cl, Br, CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 or else can carry an annelated methanediylbisoxo group or ethane-1,2-diylbisoxo group, and in addition in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl,

R^5 and $R^{5'}$, independently of one another, mean C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, whereby a carbon atom can be exchanged for O, S, SO, SO_2 , NH, N C_{1-3} alkyl or N C_{1-3} alkanoyl,

C_{3-7} cycloalkyl- C_{0-3} alkyl, whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl,

C_{6-10} aryl or 5- to 10-membered heteroaryl with 1-4 heteroatoms from N, S, and O, whereby the mentioned alkyl, alkenyl and alkynyl chains can be substituted

09759360 011601

with one of the previously mentioned cycloalkyls, aryls or heteroaryl,

whereby all previously mentioned alkyl and cycloalkyl radicals with up to two substituents consisting of CF_3 , C_2F_5 , OH , O C_{1-3} alkyl, NH_2 , NH C_{1-3} alkyl, NH C_{1-3} alkanoyl, $\text{N (C}_{1-3} \text{ alkyl)}_2$, $\text{N(C}_{1-3} \text{ alkyl)(C}_{1-3} \text{ alkanoyl)}$, COOH , CONH_2 , COO C_{1-3} alkyl and all previously mentioned aryl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F , Cl , Br , CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 or else can carry an annelated methanediylbisoxy, ethane-1,2-diylbisoxy group,

or R^5 and $\text{R}^{5'}$ together with the nitrogen atom form a 5- to 7-membered heterocyclic compound, which can contain another oxygen, nitrogen or sulfur atom and can be substituted with C_{1-4} alkyl, C_{1-4} alkoxy- C_{0-2} alkyl, C_{1-4} alkoxy-carbonyl, aminocarbonyl or phenyl,

- A means C_{1-10} alkanediyl, C_{2-10} alkenediyl, C_{2-10} alkinediyl, (C_{0-5} alkanediyl- C_{3-7} cycloalkanediyl- C_{0-5} alkanediyl), whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl, whereby in the above-mentioned aliphatic chains, a carbon atom or two carbon atoms can be exchanged for O, NH, N C_{1-3} alkyl, N C_{1-3} alkanoyl, and whereby alkyl or

09759360 011601

Sub
B1
cont

Sub
B
cont

cycloalkyl groups can be substituted with up to two substituents consisting of =O, OH, O C₁₋₃ alkyl, NH₂, NH C₁₋₃ alkyl, NH C₁₋₃ alkanoyl, N (C₁₋₃ alkyl)₂, N(C₁₋₃ alkyl) (C₁₋₃ alkanoyl),

B means COOH, COOR⁵, CONH₂, CONHNH₂, CONHR⁵, CONR⁵R^{5'}, CONHOH, CONHOR⁵, SO₃H, SO₂NH₂, SO₂NHR⁵, SO₂NR⁵R^{5'}, PO₃H, PO(OH)(OR⁵), PO(OR⁵)(OR^{5'}), PO(OH)(NHR⁵), PO(NHR⁵)(NHR^{5'}), tetrazolyl,

in each case bonded to a carbon atom of group A, or the entire group Y-A-B N(SO₂R⁴)(SO₂R^{4'}) or NHSO₂R⁴,

X means a bond, CH₂, (CH₂)₂, CH(CH₃), (CH₂)₃, CH(CH₂CH₃), CH(CH₃)CH₂, CH₂CH(CH₃),

Y means O, NH, NR⁴, NCOR⁴, NSO₂R⁴,

provided that if Y means NH, NR⁴, NCOR⁴ or NSO₂R⁴, and

a) substituent R² contains a nitrogen-containing, saturated heterocyclic compound, this heterocyclic compound is not substituted in the imine nitrogen with H, methyl, ethyl, propyl or isopropyl,

or

b) in optionally present groups XNHR⁴ or XNR⁴R^{4'} of substituent R², R⁴ and/or R^{4'} does not mean C₁₋₄ alkyl,

that B does not mean COOH, SO₃H, PO₃H₂ or tetrazolyl at the same time, and R¹ and R², independently of one another, mean C₅₋₆ heteroaryl or phenyl, if the latter, independently of one another, are unsubstituted, or are substituted simply with C₁₋₆

09759360, 011601

Sub B cont
 alkyl, C₁₋₄ perfluoroalkyl, O C₁₋₆ alkyl, O C₁₋₄ perfluoroalkyl, COOH, COO C₁₋₆ alkyl, CO C₁₋₆ alkyl, CONH₂, CONHR⁴, NO₂, NH₂, NHCOR⁴, NHSO₂R⁴, or with 1 or 2 halogen atoms from the group that consists of F, Cl, Br, and I, and whereby the following compounds are excluded:

[(1,2-Diphenyl-1H-benzimidazol-6-yl)oxy]acetic acid methyl ester,

5-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]pentanoic acid methyl ester,

4-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]butanoic acid ethyl ester,

5-[[1-(4-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

6-[[1-(4-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester,

5-[[1-(4-aminophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

5-[[1-[4-[(4-chlorophenyl)sulfonyl]amino]phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

5-[[1-[4-[(acetyl)amino]phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester

5-[[1-(3-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

6-[[1-(3-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester,

5-[[1-(3-aminophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

09759360-011601

5-[[1-[3-[[[(4-chlorophenyl)sulfonyl]amino]phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester,

5-[[1-[3-[(acetyl)amino]phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester.

2. Benzimidazoles according to claim 1, characterized in that

R¹ means a monocyclic or bicyclic C₆₋₁₂ aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-2 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br,

XOH, XOR⁴, XOCOR⁴, XOCONHR⁴, XOCOOR⁴,

XCOR⁴, XCN, XCOOH, XCOOR⁴, XCONH₂, XCONR⁴R^{4'}, XCONHR⁴,

XCONHOH, XCONHOR⁴, XCOSR⁴, XSR⁴, NO₂, XNHR⁴, XNR⁴R^{4'}, R⁴,

whereby two substituents at R¹, if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediylbisoxy, ethane-1,2-diylbisoxy, propane-1,3-diyl, butane-1,4-diyl.

3. Benzimidazoles according to claim 1 or 2, wherein

R² means a monocyclic or bicyclic C₆₋₁₀ aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-2 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or

09759360.01601
FOI TO 09E65260

heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br,

XOH, XOR⁴, XOCOR⁴, XOCONHR⁴, XOCOOR⁴,

XCOR⁴, XC(NOH)R⁴, XC(NOR⁴)R^{4'}, XC(NO(COR⁴))R^{4'},

XCOOH, XCOOR⁴, XCONH₂, XCONHR⁴, XCONR⁴R^{4'}, XCONHOH,

XCONHOR⁴, XCOSR⁴,

XSR⁴, XSOR⁴, XSO₂R⁴, SO₂NH₂, SO₂NHR⁴, SO₂NR⁴R^{4'},

NO₂, XNHR⁴, XNR⁴R^{4'}, XNHSO₂R⁴, XN(SO₂R⁴)SO₂R^{4'}, XNR⁴SO₂R^{4'},

R⁴,

whereby two substituents at R², if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediylbisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl.

4. Benzimidazoles according to one of claims 1-3, wherein R³ means one or two substituents, which, independently of one another, can be:

hydrogen, F, Cl, Br,

XOH, XOR⁴, XOCOR⁴, XOCONHR⁴, XOCOOR⁴,

XCOR⁴, XC(NOH)R⁴, XC(NOR⁴)R^{4'}, XC(NO(COR⁴))R^{4'},

XCN, XSR⁴, XSOR⁴, XSO₂R⁴, SO₂NH₂, SO₂NHR⁴, SO₂NR⁴R^{4'},

NO₂, XNH₂, XNHR⁴, XNR⁴N^{4'},

XNHSO₂R⁴, XNR⁴SO₂R^{4'}, XN(SO₂R⁴)SO₂R^{4'},

XNHCOOR⁴, XNHCONHR⁴, or R⁴, whereby two

substituents R³, if they are in ortho-position to one

FOI 0956750

another, can be linked to one another in such a way that they jointly form methanediylbisoxy, ethane-1,2-diylbisoxy, propane-1,3-diyl, or butane-1,4-diyl.

5. Benzimidazoles according to one of claims 1-4, wherein R^4 and $R^{4'}$, independently of one another, mean CF_3 , C_2F_5 , C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, (C_{1-3} alkyl- C_{3-6} cycloalkyl), phenyl or 5- to 6-membered heteroaryl with 1-2 N, S or O atoms, whereby the phenyl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F, Cl, Br, CH_3 , C_2H_5 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 ,

and in addition in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl.

6. Benzimidazoles according to one of claims 1-5, wherein R^5 and $R^{5'}$, independently of one another, can be C_{1-6} alkyl, whereby a carbon atom can be exchanged for O, NH, N C_{1-3} alkyl, N C_{1-3} alkanoyl, C_{3-7} cycloalkyl- C_{0-3} alkyl, whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl, whereby the mentioned C_{1-6} alkyl part can be substituted with one of the previously mentioned

09759360.011601
FOI 09E65460

cycloalkyls or else a 5- to 6-membered heteroaromatic compound with 1-2 heteroatoms, selected from N, S or O, whereby all previously mentioned alkyl and cycloalkyl parts can be substituted with up to two substituents that consist of CF_3 , OH, O C_{1-3} alkyl, and the previously mentioned heteroaryl groups with one or two substituents that consist of F, Cl, CF_3 , CH_3 , C_2H_5 , OCH_3 , OC_2H_5 , or R^5 and $\text{R}^{5'}$ together with the nitrogen atom form a 5- to 7-membered heterocyclic compound, which can contain another oxygen, nitrogen or sulfur atom and can be substituted with C_{1-4} alkyl, C_{1-4} alkoxy- C_{0-2} alkyl, C_{1-4} alkoxy-carbonyl, aminocarbonyl or phenyl.

7. Benzimidazoles according to one of claims 1-6, wherein
- A means C_{1-10} alkanediyl, C_{2-10} alkenediyl, C_{2-10} alkinediyl, (C_{0-5} alkanediyl- C_{3-7} cycloalkanediyl- C_{0-5} alkanediyl), whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O, or in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl, whereby in the above-mentioned aliphatic chains, a carbon atom or two carbon atoms can be exchanged for O, NH, N C_{1-3} alkyl, or N C_{1-3} alkanoyl.
8. Benzimidazoles according to one of claims 1-7, wherein
- B means COOH , COOR^5 , CONH_2 , CONHR^5 , $\text{CONR}^5\text{R}^{5'}$, CONHOH , CONHOR^5 or tetrazolyl, in each case bonded to a carbon atom of group A.

09750360-011601

9. Benzimidazoles according to one of claims 1-8, wherein

X means a bond or methylene.

10. Benzimidazoles according to one of claims 1-9, wherein

Y means O.

11. [(1,2-Diphenyl-1H-benzimidazol-6-yl)oxy]acetic acid

isopropyl ester

3-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]propanoic acid

methyl ester

2-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]propanoic acid

methyl ester

4-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]butanoic acid

isopropyl ester

5-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]pentanoic acid

isopropyl ester

6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanoic acid

methyl ester

6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanoic acid

isopropyl ester

6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-methoxy-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-(phenylmethoxy)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-hydroxy-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

7-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]heptanoic acid
methyl ester

009759360-011601

6-[[1-(3-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[2-phenyl-1-[3-(trifluoromethyl)phenyl]-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-[3-(trifluoromethyl)phenyl]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3-cyanophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(3-cyanophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3-cyanophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[1-(4-cyanophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-cyanophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3-chlorophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(3-chlorophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(4-chlorophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-chlorophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

00759360 011601

6-[[1-(3-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(3,5-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(3,5-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-(3-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(3,4-dimethoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-[3,4-(methylenedioxy)phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-[3,4-(methylenedioxy)phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid

09759360-011601

6-[[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-[4-(N,N-dimethylamino)phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-[4-(N,N-dimethylamino)phenyl]-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[1-phenyl-2-[3-(trifluoromethyl)phenyl]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[2-(3-chlorophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(3-chlorophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[2-(4-chlorophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-chlorophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[2-(4-methylphenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-methylphenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1-phenyl-2-(4-pyridinyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[(1,2-diphenyl-5-nitro-1H-benzimidazol-6-yl)oxy]hexanoic acid methyl ester

6-[(1,2-diphenyl-5-nitro-1H-benzimidazol-6-yl)oxy]hexanoic acid isopropyl ester

09759360-011601

Sub
C4
Cont.

6-[[5-[[[4-bromophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[5-[[[4-chlorophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[[4-chlorophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1,2-diphenyl-5-[[[3-methylphenyl)sulfonyl]amino]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1,2-diphenyl-5-[[[4-methylphenyl)sulfonyl]amino]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1,2-diphenyl-5-[[[4-methoxyphenyl)sulfonyl]amino]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1,2-diphenyl-5-[[[4-trifluoromethyl)phenyl)sulfonyl]amino]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[5-[[[4-(acetylamino)phenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]-hexanoic acid isopropyl ester

6-[[5-[[bis(3-chlorophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[1,2-diphenyl-5-[(propylsulfonyl)amino]-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[5-[(benzylsulfonyl)amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

2-[2-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]ethoxy]acetic acid methyl ester

3-[2-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]ethoxy]propanoic acid methyl ester

00759360-011601

Sub
C4
cont.

6-[[1-(3-nitrophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid ethyl ester

6-[[4-acetyl-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-5-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-[4-(thiomethyl)phenyl]-1H-benzimidazol-5-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-[4-(thiomethyl)phenyl]-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-(3-thienyl)-1H-benzimidazol-5-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-(3-thienyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

4-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]butanoic acid methyl ester

N-(phenylmethoxy)-6-[[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]-hexanamide

N,N-dimethyl-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-isopropyl-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]-1-pyrrolidin-1-ylhexan-1-one

5-[[5-[[4-chlorophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester

Sub
C4
cont.

009360-011601

6-[[5-[[[(4-chlorophenyl)sulfonyl]amino]-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[[(4-chlorophenyl)sulfonyl]amino]-1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[4-(acetyloxy)-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[4-hydroxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[4-hydroxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[7-methyl-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

12. 6-[[2-Phenyl-1-(3-pyridyl)-1H-benzimidazol-5-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-(3-pyridyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-phenyl-1-(4-pyridyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-fluoro-phenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-methoxyphenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-bromophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-[4-(trifluoromethyl)phenyl]-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

0911601-011601

Sub
C4
cont.

6-[[1-phenyl-2-(benzothien-2-yl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-phenyl-2-(benzothien-2-yl)-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[5-hydroxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[5-hydroxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[5-methoxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid isopropyl ester

6-[[5-hydroxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-methoxy-1-(4-methylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[4-chlorophenyl]sulfonyl]amino]-1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[4-chlorophenyl]sulfonyl]amino]-2-(4-fluorophenyl)-1-(4-methoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[4-chlorophenyl]sulfonyl]amino]-1-(4-methoxyphenyl)-2-(4-methoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

4-[[5-[[4-chlorophenyl]sulfonyl]amino]-1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]butanoic acid methyl ester

5-[[5-[[4-chlorophenyl]sulfonyl]amino]-1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester

003360 011601

5-[[5-[[[(4-chlorophenyl)sulfonyl]amino]-1,2-diphenyl-1H-benzimidazol-6-yl]oxy]pentanoic acid methyl ester

6-[[5-[[[(4-(trifluoromethyl)phenyl)sulfonyl]amino]-1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[5-[[[(4-chlorophenyl)sulfonyl]methylamino]-1-(4-methoxyphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(indan-5-yl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(indan-5-yl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid

6-[[1-(3-fluorophenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(4-nitrophenyl)-1-phenyl-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-phenyl-2-(3-pyridinyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

N-(cyclopropylmethoxy)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-isobutoxy-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-(cyclopropylmethoxy)-6-[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]-hexanamide

N-isobutoxy-6-[2-phenyl-1-(3,4,5-trimethoxyphenyl)-1H-benzimidazol-6-yl]oxy]hexanamide

04759360.014601

N-(2-methoxyethyl)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-(3-methoxypropyl)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-isobutyl-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]-1-morpholin-1-ylhexan-1-one

N,N-di(-2-methoxyethyl)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-isopentyl-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-(pyridin-2-yl)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-(pyridin-3-yl)-6-[(1,2-diphenyl-1H-benzimidazol-6-yl)oxy]hexanamide

N-isopropyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N,N-dimethyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N,N-diethyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N-isobutyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N-cyclopropyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

059360.01601

N-cyclobutyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N-tert-butyl-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

(R)-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]1-(2-methoxymethyl)-pyrrolidin-1-ylhexan-1-one

N-(3-imidazol-1-yl-propyl)-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N-(2-pyridin-2-ylethyl)-6-[[1-(3,4-dimethylphenyl)-2-phenyl-1H-benzimidazol-6-yl]oxy]hexanamide

N-(3-methoxypropyl)-6-[[1-(indan-5-yl)-2-phenyl-1H-benzimidazol-6-yl]oxy]heptanamide

6-[[1-(4-methylphenyl)-2-(3-pyridyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(4-pyridyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(2-thienyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(3-thienyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[2-(3-indolyl)-1-(4-methylphenyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(2-furyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(3-furyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

19750360-011601


6-[[1-(4-methylphenyl)-2-(5-methyl-2-thienyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester

6-[[1-(4-methylphenyl)-2-(3-methyl-2-thienyl)-1H-benzimidazol-6-yl]oxy]hexanoic acid methyl ester.

13. Use of a compound according to one of claims 1-12 for the production of a pharmaceutical agent for treating or preventing diseases that are associated with a microglia activation.

14. Pharmaceutical agent, wherein it contains one or more compounds according to one of claims 1-12 and one or more vehicles.

15. Use of a benzimidazole of general formula II



in which

R^1 means a monocyclic or bicyclic C_{6-12} aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-4 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

09759360.014604

Sub
B2
cont

Sub
B₂
cont

F, Cl, Br, I, C(NH)NH₂, C(NH)NHR⁴, C(NH)NR⁴R^{4'},
C(NR⁴)NH₂, C(NR⁴)NHR^{4'}, C(NR⁴)NR⁴R^{4'}, XOH, XOR⁴, XOCOR⁴,
XOCONHR⁴, XOCOOR⁴, XCOR⁴, XC(NOH)R⁴, XC(NOR⁴)R^{4'},
XC(NO(COR⁴))R^{4'}, XCN, XCOOH, XCOOR⁴, XCONH₂, XCONR⁴R^{4'},
XCONHR⁴, XCONHOH, XCONHOR⁴, XCOSR⁴, XSR⁴, XSOR⁴, XSO₂R⁴,
SO₂NH₂, SO₂NHR⁴, SO₂NR⁴R^{4'}, NO₂, XNH₂, XNHR⁴, XNR⁴R^{4'},
XNH₂SO₂R⁴, XN(SO₂R⁴)(SO₂R^{4'}), XNR⁴SO₂R^{4'}, XNHCOR⁴, XNHCOOR⁴,
XNHCONHR⁴, tetrahydro-2,5-dioxopyrrol-1-yl, 2,5-
dihydro-2,5-dioxopyrrol-1-yl, 2,7-dihydro-2,7-
dioxoisindol-1-yl, R⁴, whereby two substituents at R¹,
if they are in ortho-position to one another, can be
linked to one another in such a way that they jointly
form methanediylbisoxo, ethane-1,2-diylbisoxo, propane-
1,3-diyl, butane-1,4-diyl,

R² means a monocyclic or bicyclic C₆₋₁₀ aryl group or a
monocyclic or bicyclic 5- to 10-membered heteroaryl
group with 1-4 heteroatoms selected from the group that
consists of N, S or O, whereby the mentioned aryl or
heteroaryl group can be substituted with up to three of
the following substituents, independently of one
another:

F, Cl, Br, I, C(NH)NH₂, C(NH)NHR⁴, C(NH)NR⁴R^{4'},
C(NR⁴)NH₂, C(NR⁴)NHR^{4'}, C(NR⁴)NR⁴R^{4'}, XOH, XOR⁴, XOCOR⁴,
XOCONHR⁴, XOCOOR⁴, XCOR⁴, XC(NOH)R⁴, XC(NOR⁴)R^{4'},
XC(NO(COR⁴))R^{4'}, XCN, XCOOH, XCOOR⁴, XCONH₂, XCONR⁴R^{4'},
XCONHR⁴, XCONHOH, XCONHOR⁴, XCOSR⁴, XSR⁴, XSOR⁴, XSO₂R⁴,
SO₂NH₂, SO₂NHR⁴, SO₂NR⁴R^{4'}, NO₂, XNH₂, XNHR⁴, XNR⁴R^{4'},

Sub
B2
cont

$\text{XNH}\text{SO}_2\text{R}^4$, $\text{XN}(\text{SO}_2\text{R}^4)(\text{SO}_2\text{R}^{4'})$, $\text{XNR}^4\text{SO}_2\text{R}^{4'}$, XNHCOR^4 , XNHCOOR^4 , XNHCONHR^4 , tetrahydro-2,5-dioxopyrrol-1-yl, 2,5-dihydro-2,5-dioxopyrrol-1-yl, 2,7-dihydro-2,7-dioxoisindol-1-yl, R^4 , whereby two substituents at R^2 , if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediyl-bisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl,

R^3 stands for one or two substituents, which form, independently of one another:

hydrogen, F, Cl, Br, I, XOH , XOR^4 , XOCOR^4 , XOCONHR^4 , XOCOOR^4 , XCOR^4 , $\text{XC}(\text{NOH})\text{R}^4$, $\text{XC}(\text{NOR}^4)\text{R}^{4'}$, $\text{XC}(\text{NO}(\text{COR}^4))\text{R}^{4'}$, XCN , XCOOH , XCOOR^4 , XCONH_2 , XCONHR^4 , $\text{XCONR}^4\text{R}^{4'}$, XCONHOH , XCONHOR^4 , XCOSR^4 , XSR^4 , XSOR^4 , XSO_2R^4 , SO_2NH_2 , SO_2NHR^4 , $\text{SO}_2\text{NR}^4\text{R}^{4'}$, NO_2 , XNH_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$, $\text{XNH}\text{SO}_2\text{R}^4$, $\text{XNR}^4\text{SO}_2\text{R}^{4'}$, $\text{XN}(\text{SO}_2\text{R}^4)(\text{SO}_2\text{R}^{4'})$, XNHCOR^4 , XNHCOOR^4 , XNHCONHR^4 , tetrahydro-2,5-dioxopyrrol-1-yl, 2,5-dihydro-2,5-dioxopyrrol-1-yl, 2,7-dihydro-2,7-dioxoisindol-1-yl, R^4 , whereby two substituents at R^3 , if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediylbisoxo, ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-diyl,

R^4 and $\text{R}^{4'}$, independently of one another, mean C_{1-4}

perfluoroalkyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_3 , 7 cycloalkyl, (C_{1-3} alkyl- C_{3-7} cycloalkyl), C_{1-3} alkyl- C_{6-10} aryl, C_{1-3} alkyl 5 to 10-membered heteroaryl, with 1-4

09759360-011601

Sub
B2
cont

N, S or O atoms, C_{6-10} aryl or 5- to 10-membered heteroaryl with 1-4 N, S or O atoms, whereby the C_{6-10} aryl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F, Cl, Br, CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 or else can carry an annelated methanediylbisoxo group or ethane-1,2-diylbisoxo group, and in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl,

R^5 and $R^{5'}$, independently of one another, mean hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, whereby a carbon atom can be exchanged for O, S, SO, SO_2 , NH, N C_{1-3} alkyl or N C_{1-3} alkanoyl,

C_{3-7} cycloalkyl- C_{0-3} alkyl, whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl,

C_{6-10} aryl or 5- to 10-membered heteroaryl with 1-4 heteroatoms from N, S, and O, whereby the mentioned alkyl, alkenyl and alkynyl chains can be substituted with one of the previously mentioned cycloalkyls, aryls or heteroaryls,

09759360-011601

Sub
B2
ent

whereby all previously mentioned alkyl and cycloalkyl radicals with up to two substituents consisting of CF_3 , C_2F_5 , OH , O C_{1-3} alkyl, NH_2 , NH C_{1-3} alkyl, NH C_{1-3} alkanoyl, $\text{N (C}_{1-3} \text{ alkyl)}_2$, $\text{N(C}_{1-3} \text{ alkyl)(C}_{1-3} \text{ alkanoyl)}$, COOH , CONH_2 , COO C_{1-3} alkyl and all previously mentioned aryl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F , Cl , Br , CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 or else can carry an annelated methanediylbisoxy, ethane-1,2-diylbisoxy group, or R^5 and $\text{R}^{5'}$ together with the nitrogen atom form a 5-to 7-membered heterocyclic compound, which can contain another oxygen, nitrogen or sulfur atom and can be substituted with C_{1-4} alkyl, C_{1-4} alkoxy- C_{0-2} alkyl, C_{1-4} alkoxy-carbonyl, aminocarbonyl or phenyl,

- A means C_{1-10} alkanediyl, C_{2-10} alkenediyl, C_{2-10} alkinediyl, $(\text{C}_{0-5} \text{ alkanediyl-C}_{3-7} \text{ cycloalkanediyl-C}_{0-5} \text{ alkanediyl})$, $(\text{C}_{0-5} \text{ alkanediylarylene-C}_{0-5} \text{ alkanediyl})$, $(\text{C}_{0-5} \text{ alkanediyl-heteroarylene-C}_{0-5} \text{ alkanediyl})$, whereby the aryl and heteroaryl groups can be substituted with one or two substituents that consist of F , Cl , Br , CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 , whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O , and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O , whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl,

09759360-011601

Sub B2 cont

whereby in the mentioned aliphatic chains, a carbon atom or two carbon atoms can be exchanged for O, NH, NR^4 , NCOR^4 , NSO_2R^4 , and whereby alkyl or cycloalkyl groups can be substituted with up to two substituents consisting of F, OH, OR^4 , OCOR^4 , =O, NH_2 , $\text{NR}^4\text{R}^{4'}$, NHCOR^4 , NHCOOR^4 , NHCONHR^4 , NHSO_2R^4 , SH, SR^4 ,

B means hydrogen, OH, OCOR^5 , OCONHR^5 , OCOOR^5 , COR^5 , $\text{C}(\text{NOH})\text{R}^5$, $\text{C}(\text{NOR}^5)\text{R}^{5'}$, $\text{C}(\text{NO}(\text{COR}^5))\text{R}^{5'}$, COOH , COOR^5 , CONH_2 , CONHNH_2 , CONHR^5 , $\text{CONR}^5\text{R}^{5'}$, CONHOH , CONHOR^5 , SO_3H , SO_2NH_2 , SO_2NHR^5 , $\text{SO}_2\text{NR}^5\text{R}^{5'}$, PO_3H , $\text{PO}(\text{OH})(\text{OR}^5)$, $\text{PO}(\text{OR}^5)(\text{OR}^{5'})$, $\text{PO}(\text{OH})(\text{NHR}^5)$, $\text{PO}(\text{NHR}^5)(\text{NHR}^{5'})$, tetrazolyl, respectively bonded to a carbon atom of group A,

or the entire group $\text{Y-A-B N}(\text{SO}_2\text{R}^4)(\text{SO}_2\text{R}^{4'})$ or NHSO_2R^4 ,

X means a bond, CH_2 , $(\text{CH}_2)_2$, $\text{CH}(\text{CH}_3)$, $(\text{CH}_2)_3$, $\text{CH}(\text{CH}_2\text{CH}_3)$, $\text{CH}(\text{CH}_3)\text{CH}_2$, $\text{CH}_2\text{CH}(\text{CH}_3)$,

Y means a bond, O, S, SO, SO_2 , NH, NR^4 , NCOR^4 , NSO_2R^4 , for the production of a pharmaceutical agent for treating or preventing diseases that are associated with a microglia activation.

16. Use according to claim 15, whereby in general formula II,

R^1 means a monocyclic or bicyclic aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl

09759360-011601

Sub B2 cont
group with 1-2 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br,

XOH, XOR^4 , XOCOR^4 , XOCONHR^4 , XOCOOR^4 ,

XCOR^4 , XCN , XCOOH , XCOOR^4 , XCONH_2 , $\text{XCONR}^4\text{R}^{4'}$, XCONHR^4 ,

XCONHOH ,

XCONHOR^4 , XCOSR^4 , XSR^4 , NO_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$,

R^4 ,

whereby two substituents at R^1 , if they are in ortho-position to one another, can be linked to one another in such a way that they jointly form methanediylbisoxy, ethane-1,2-diylbisoxy, propane-1,3-diyl, butane-1,4-diyl.

17. Use according to claim 15 or 16, whereby in general formula II,

Sub A3
 R^2 means a monocyclic or bicyclic aryl group or a monocyclic or bicyclic 5- to 10-membered heteroaryl group with 1-2 heteroatoms selected from the group that consists of N, S or O, whereby the mentioned aryl group or heteroaryl group can be substituted with up to three of the following substituents, independently of one another:

F, Cl, Br, XOH, XOR^4 , XOCOR^4 , XOCONHR^4 , XOCOOR^4 ,

XCOR^4 , $\text{XC}(\text{NOH})\text{R}^4$,

09759360-011604

$\text{XC}(\text{NOR}^4)\text{R}^{4'}$, $\text{XC}(\text{NO}(\text{COR}^4))\text{R}^{4'}$, XCN , XCOOH , XCOOR^4 , XCONH_2 ,
 $\text{XCONR}^4\text{R}^{4'}$,
 XCONHR^4 , XCONHOH , XCONHOR^4 , XCOSR^4 , XSR^4 , XSOR^4 , XSO_2R^4 ,
 SO_2NH_2 , SO_2NHR^4 , $\text{SO}_2\text{NR}^4\text{R}^{4'}$, NO_2 , XNH_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$,
 $\text{XNH}\text{SO}_2\text{R}^4$,
 $\text{XN}(\text{SO}_2\text{R}^4)(\text{SO}_2\text{R}^{4'})$, $\text{XNR}^4\text{SO}_2\text{R}^{4'}$, XNHCOR^4 , XNHCOOR^4 ,
 XNHCONHR^4 , R^4 ,

whereby two substituents at R^2 , if they are in ortho-
 position to one another, can be linked to one another
 in such a way that they jointly form methanediylbisoxo,
 ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-
 diyl.

18. Use according to claims 15-17, whereby in general
 formula II

R^3 stands for one or two substituents, which independently
 of one another, mean:

hydrogen, F, Cl, Br, XOH , XOR^4 , XOCOR^4 , XOCONHR^4 ,
 XOCOOR^4 , XCOR^4 , $\text{XC}(\text{NOH})\text{R}^4$, $\text{XC}(\text{NOR}^4)\text{R}^{4'}$, $\text{XC}(\text{NO}(\text{COR}^4))\text{R}^{4'}$,
 XCN , XSR^4 , XSOR^4 , XSO_2R^4 , SO_2NH_2 , SO_2NHR^4 , $\text{SO}_2\text{NR}^4\text{R}^{4'}$, NO_2 ,
 XNH_2 , XNHR^4 , $\text{XNR}^4\text{R}^{4'}$, $\text{XNH}\text{SO}_2\text{R}^4$, $\text{XNR}^4\text{SO}_2\text{R}^{4'}$,
 $\text{XN}(\text{SO}_2\text{R}^4)(\text{SO}_2\text{R}^{4'})$, XNHCOR^4 , XNHCOOR^4 , XNHCONHR^4 , or R^4 ,

whereby two substituents R^3 , if they are in ortho-
 position to one another, can be linked to one another
 in such a way that they jointly form methanediylbisoxo,
 ethane-1,2-diylbisoxo, propane-1,3-diyl, butane-1,4-
 diyl.

09759360-011601

19. Use according to claims 15-18, whereby in general formula II

13
 R^4 and $R^{4'}$, independently of one another, mean CF_3 , C_2F_5 , C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, (C_{1-3} alkyl- C_{3-6} cycloalkyl), C_{1-3} alkylaryl, C_{1-3} alkylheteroaryl, monocyclic aryl or 5- to 6-membered heteroaryl with 1-2 N, S or O atoms, whereby the aryl and heteroaryl groups can be substituted with one or two substituents from the group that consists of F, Cl, Br, CH_3 , C_2H_5 , NO_2 , OCH_3 , OC_2H_5 , CF_3 , C_2F_5 or else can carry an annelated methanediylbisoxy or ethane-1,2-diylbisoxy group, and in addition in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl.

20. Use according to claims 15-19, whereby in general formula II

R^5 and $R^{5'}$, independently of one another, can be C_{1-6} alkyl, whereby a carbon atom can be exchanged for O, NH, N C_{1-3} alkyl, N C_{1-3} alkanoyl, C_{3-7} cycloalkyl- C_{0-3} alkyl, whereby in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N and/or O, whereby ring nitrogens optionally can be substituted with C_{1-3} alkyl or C_{1-3} alkanoyl, whereby the

09759360.011604

mentioned C₁₋₆ alkyl part can be substituted with one of the previously mentioned cycloalkyls or else a 5- to 6-membered heteroaromatic compound with 1-2 heteroatoms, selected from the group that consists of N, S or O, whereby all previously mentioned alkyl and cycloalkyl parts can be substituted with up to two substituents that consist of CF₃, OH, O C₁₋₃ alkyl, and the previously mentioned heteroaryl groups can be substituted with one or two substituents that consist of F, Cl, CF₃, CH₃, C₂H₅, OCH₃, OC₂H₅, or R⁵ and R^{5'} together with the nitrogen atom form a 5- to 7-membered heterocyclic compound, which can contain another oxygen, nitrogen or sulfur atom and can be substituted with C₁₋₄ alkyl, C₁₋₄ alkoxy-C₀₋₂ alkyl, C₁₋₄ alkoxy-carbonyl, aminocarbonyl or phenyl.

21. Use according to claims 15-20, whereby in general formula II

A means C₁₋₁₀ alkanediyl, C₂₋₁₀ alkenediyl, C₂₋₁₀ alkinediyl, (C₀₋₅ alkanediyl-C₃₋₇ cycloalkanediyl-C₀₋₅ alkanediyl), or (C₀₋₅ alkanediyl-heteroarylene-C₀₋₅ alkanediyl), whereby an optionally present heteroaryl group can be substituted with one or two substituents that consist of F, Cl, Br, CH₃, C₂H₅, NO₂, OCH₃, OC₂H₅, CF₃, C₂F₅, and in addition in a 5-membered cycloalkyl ring, a ring member can be an N or an O, and in a 6- or 7-membered cycloalkyl ring, one or two ring members can be N

09759360-011601

and/or O, whereby ring nitrogens optionally can be substituted with C₁₋₃ alkyl or C₁₋₃ alkanoyl, whereby in an aliphatic chain, a carbon atom or two carbon atoms can be exchanged for O, NH, N C₁₋₃ alkyl, N C₁₋₃ alkanoyl, NSO₂ C₁₋₃ alkyl, and whereby alkyl or cycloalkyl parts can be substituted with up to two F atoms or one of the substituents that consists of OH, O C₁₋₃ alkyl, O C₁₋₃ alkanoyl, =O, NH₂, NH C₁₋₃ alkyl, N (C₁₋₃ alkyl)₂, NH C₁₋₃ alkanoyl, N (C₁₋₃ alkyl) (C₁₋₃ alkanoyl), NHCOO C₁₋₃ alkyl, NHCONH C₁₋₃ alkyl, NHSO₂ C₁₋₃ alkyl, SH, S C₁₋₃ alkyl.

22. Use according to claims 15-21, whereby in general formula II

B means hydrogen, OH, OCOR⁵, OCONHR⁵, OCOOR⁵, COOH, COOR⁵, CONH₂, CONHR⁵, CONR⁵R^{5'}, CONHOH, CONHOR⁵, or tetrazolyl, in each case bonded to a carbon atom of group A.

23. Use according to claims 15-22, whereby in general formula II,

X means a bond or CH₂.

24. Use according to claims 15-23, whereby in general formula II,

Y means a bond, O, S, NH, NR⁴, NCOR⁴ or NSO₂R⁴.

09759360-011601

Add B3

add C8